

On the Local Finite Density Relativistic Quantum Field Theories

S. Ying

Physics Department, Fudan University, Shanghai 200433, People's Republic of China

It is found that, in addition to the conventional ones, a local approach to the relativistic quantum field theories at both zero and finite density consistent with the violation of Bell like inequalities should contain, and provide solutions to at least three additional problems, namely, 1) the statistical gauge invariance 2) the dark components of the local observables and 3) the fermion statistical blocking effects, base upon an asymptotic non-thermo ensemble. An application to models are presented to show the importance of the discussions.

PACS Number: 11.90.+t, 11.15.Tk, 12.40.-y, 12.90.+b, 03.65.Bz

Besides its relevance to domestic processes like in heavy ion collisions, in nuclear matter, a consistent theory for relativistic finite density systems is neither less interesting fundamentally due to its relevance to profound cosmological questions like the mechanism for baryogenesis, the nature of dark matter, etc., nor can it be expected to be trivially derivable from the theoretical framework at zero density. Although such kind of theory, which adopt the basic framework of the field theoretical representation of non-relativistic many body systems exists and is widely used in literature, one essential ingredient of the relativistic spacetime, namely the principle of locality, has not been properly addressed. This is because in the grand canonical ensemble (GCE) base on which such a theory is built, the chemical potential is a global quantity in spacetime. As it is well know in the studies of few and many body relativistic systems that it is not enough to simply adopt the relativistic kinematics for each particles in the system when the interaction is present. The less well understood relative time between particles inside the system exists due to the lack of a frame independent definition of simultaneity in the relativistic spacetime which constitutes another fundamental difference between non-relativistic systems based upon Newtonian spacetime and the relativistic ones.

A measurement of the ground state fermion number density in GCE given by

$$\bar{n} = \lim_{T \rightarrow 0, \Omega \rightarrow \infty} \frac{T}{\Omega} \frac{\partial \ln Z}{\partial \mu} \quad (1)$$

corresponds to a global measurement. Here Z is the partition function, T is the temperature, μ is the chemical potential and Ω is the volume of the system. Global measurements are not directly definable since simultaneity of two events separated from each other by a space-like distance depends on reference frame in relativity. In addition, the property of asymptotic freedom in strong interaction makes the local measurements relevant to the study of the properties of fundamental current quarks at short distances where they are point like and almost free particles.

Local measurements in quantum field theory are realized by exerting an external local field to the system at the spacetime point interested, and, the measured quantity are deduced from the response of the system to the external field. For finite density systems, a Lorentz 4-vector local field $\mu^\alpha(x)$, called the primary statistical gauge field, is introduced. The ground state expectation value of fermion number density can be expressed as the functional derivative

$$\bar{\rho}(x) = \frac{\delta \ln Z}{\delta \mu^0(x)} \quad (2)$$

with Z a functional of μ^α . It corresponds to a local measurement in the ground state at the spacetime point x .

That $\bar{\rho} \equiv \bar{n}$ is not mathematically warranted. Because $\bar{\rho}$ contains not only the contributions from the quasiparticles but also the ones from localized transient fluctuations of the fields [1] while $\bar{n}$ contains only the quasiparticle contributions. The difference $\rho_D = \bar{\rho} - \bar{n}$ is called the dark component of the local observable.

$\bar{\rho}$ follows from the asymptotic grand canonical ensemble (AGCE), which is defined as the GCE of the corresponding free system in the remote past (or future) when the interaction effects are adiabatically switched off, with the Lagrangian density that underlies Z given by

$$\mathcal{L}' = \frac{1}{2} \bar{\Psi} (i \not{\partial} + \not{O}_3 - m) \Psi + \mathcal{L}_B + \mathcal{L}_{\text{int}} \quad (3)$$

and m the fermion mass, $\mathcal{L}_B$ the Lagrangian density of boson fields of the system and $\mathcal{L}_{\text{int}}$ the interaction between the boson and fermion fields. An 8 component “real” fermion field Ψ is used for the discussion [2] with O_3 the third Pauli matrices acting on the upper and lower 4-component of Ψ . In most of the cases, $\mathcal{L}_{\text{int}} = -\bar{\Psi}(\Sigma(f) - m)\Psi$ with

Σ a function(al) of the boson fields represented by f so that the fermion degrees of freedom can be integrated out leading to an effective Euclidean action for the boson fields

$$S_{\text{eff}}[f, \mu] = \frac{1}{2} \text{SpLn} \frac{[i\partial + \not{m} O_3 - \Sigma(f)]}{[i\partial + \not{m} O_3 - m]} + \int d^4x [\mathcal{L}_B(f) + \mu \bar{\rho} - \bar{e}] \quad (4)$$

with “Sp” the functional trace, $\bar{e}$ the average energy density of the corresponding free system of fermions ($\Sigma = 0$) and $\bar{\rho}$ is the local average of fermion number density to be discussed in the following at the “chemical potential” $\mu \equiv \sqrt{\bar{\mu}^\alpha \bar{\mu}_\alpha}$, where $\bar{\mu}^\alpha$ is the ground state value of μ^α normally provided by the external conditions. The partition functional $Z = \int D[f] \exp(S_{\text{eff}})$ is then a functional of μ^α and $\ln Z$ its effective action.

The first term in Eq. 4 is invariant under the following $U(1)$ statistical gauge transformation $\mu^\alpha(x) \rightarrow \mu^\alpha(x) - \partial^\alpha \Lambda(x)$, which corresponds to the conservation of the fermion number. Since μ^α is a local field, its excitation represents certain collective excitations of the system. Such a view introduces infinite many extra degrees of freedom since there is no such a field in the original theory. Superselection-sector in the representing Hilbert space (SIRHS) containing physical states correspond to the primary statistical gauge field exists and can be selected using a “Gauss Law” constraint. For the statistical gauge invariant system, it can be imposed differently on the physical eigenstates of the Hamiltonian without causing contradictions. States in a SIRHS identified by a coordinate dependent complex function ς satisfy

$$\langle \psi_\varsigma^i | (\hat{\rho} + \nabla \cdot \hat{\pi}_u) | \psi_\varsigma^j \rangle = \delta_{E_i E_j} N_{ij} \varsigma \quad (5)$$

with $\hat{\pi}_u$ the “statistical electric field”, $|\psi_\varsigma^k\rangle$ a physical state that has energy E_k , $\delta_{EE'}$ taking zero or unity value if $E \neq E'$ or $E' = E$ (assuming that E is discrete before thermodynamic limit is taken) and N_{ij} independent of spacetime. $\hat{\rho} + \nabla \cdot \hat{\pi}_u$ is the generator of local statistical gauge transformations, so the physical states in a SIRHS change a common (coordinate dependent) phase under a specific gauge transformation rather than remains invariant, which has hitherto been used. Clearly, the later condition is a special case for the former.

Due to the conservation of the fermion number, the SIRHS of the system with fixed fermion number selected [3] by the AGCE is invariant during the time evolution. In the AGCE, the dependence of $\bar{\rho}$ upon the spacetime independent μ remains the same whether there is interaction in the system or not. It is given by $\bar{\rho} = N_g \mu^3 / 3\pi^2$ for a massless system, with N_g the total internal degrees of freedom besides the spin of the fermion. $\bar{\rho}$ does not has a simple dependence on μ , especially when there is certain kind of phase transition that modifies the excitation spectra of the system.

In some sense, the AGCE is a GCE for the initial state of the system under the time evolution, it is a canonical ensemble for the interaction effects since the invariant SIRHS (under the time evolution) is fixed in the remote past (or future). The usefulness of the AGCE in numerical simulations has already manifested in Monte-Carlo studies of the finite density problems [4] based upon GCE for a finite lattice. It is shown using the chiral Gross-Neveu model that for a given chemical potential, the fermion number density of the system has two stable values, which are reached by generating the Monte-Carlo ensemble at either finite or zero density. The first stable one correspond to the one given by the AGCE $\rho - \mu$ relation and the second one correspond to the GCE prediction for a massive system of quasi-particles. The exact results for the global treatment of the problem crossover from the later $\rho - \mu$ curve to the former one near the chiral symmetry restoration $\mu = \mu_c$. This behavior can be interpreted as that before the chiral symmetry restoration, the first $\rho - \mu$ curve for the AGCE is a local minimum of the (Euclidean) action of the various configurations in the Monte-Carlo ensemble with the GCE curve for the massive system of quasiparticles the absolute one. Near the chiral symmetry restoration point, their role exchanges. The existence of a stable local minimum for the AGCE $\rho - \mu$ curve constitutes a strong numerical evidence that the AGCE is useful in numerical simulations of the localized theories. In fact, it is expected that some of the puzzles encountered in the lattice gauge theory studies at finite density, like the early on set of the baryon number before the chiral symmetry restoration, can be understood in the light of the AGCE since the finite size effects of the lattice gauge theory study based upon GCE corresponds to, to some degree, a low resolution local measurement of an infinite system [1,2].

This interpretation can be substantiated by using a cluster decomposition approximation of the partition functional $Z[\mu]$ [1] which is expected to provide a non-perturbative picture for the μ dependence of the fermion number density at sufficiently low spacetime resolutions. In the crudest approximation, the ground state fermion number density can be written as [1]

$$\rho_{\Delta\Omega}(\bar{\sigma}, \mu) = \sqrt{\frac{\alpha}{\pi}} \int_{-\mu-\bar{\sigma}}^{\mu-\bar{\sigma}} d\sigma' e^{-\alpha\sigma'^2} \rho_\infty(\bar{\sigma} + \sigma', \mu), \quad (6)$$

where $\bar{\sigma}$ is the ground state expectation value of σ , $\rho_\infty(\sigma, \mu) = g(\mu^2 - \sigma^2)^{3/2}/3\pi^2$ is the ground state fermion number density in the GCE with fermion mass σ . Here $\Delta\Omega$ is the spacetime volume that can be resolved in the observation. α is proportional to $\Delta\Omega$ for large enough $\Delta\Omega$ and approaches a constant value for small $\Delta\Omega$ which scales with the inverse mass gap of the σ' excitation of the system. Clearly, $\rho_{\Delta\Omega}(\bar{\sigma}, \mu) \rightarrow \rho_\infty(\bar{\sigma}, \mu) = \bar{n}$ as $\Delta\Omega \rightarrow \infty$. This qualitative picture shows that the result of a local measurement of the fermion number density approaches that of the global one as the resolution gets lowered; in the limit of the zero resolution, the result approaches that of the global measurement as expected.

The existence of the dark component of local fermion number density has interesting implications. Take the vacuum state of a massless strongly interacting system in which the chiral symmetry is spontaneously broken down for example. The originally gapless vacuum state acquires an excitation gap. If only the massive quasi-particles are taken into account, the baryon number density are expected to be non-vanishing only when μ is larger than the mass of the quasi-particles. The local fermion number density is however finite as long as μ is not zero. A natural question arises as to what this dark component of fermion number corresponding to? It is reasonable to conjecture that this dark component of fermion number corresponds to those fermion states that are localized and non-propagating [5] similar to the Anderson localization in a condensed matter system. The random potential here is self-generated within the system by the transient and localized random quantum fluctuations of the σ and other boson fields. Such a conjecture is amenable to future studies.

At the fundamental level, the dark component of local observables is of pure quantum in origin related to the space-like correlations between local measurements in the relativistic quantum world that are shown to exist in experimental observations [6–8], which are what reality manifests itself [9]. This is because the size of the correlated cluster used in studying the vacuum state (in the Euclidean spacetime) tends to zero in the classical limit ($\hbar \rightarrow 0$) as a result of the classical relativistic causality, which suppresses the dark component for any finite resolution observation [1].

The Euclidean effective action for the boson field given by Eq. 4 can be evaluated in the usual way [2]. Since the effective action given by Eq. 4 is a canonical functional of μ^α , we can make a Legendre transformation of it, namely $\tilde{S}_{\text{eff}} = S_{\text{eff}} - \int d^4x \bar{\mu}_\alpha \bar{j}^\alpha$ with $\bar{j}^\alpha$ the ground state current of fermion number density, to make it a canonical functional of the fermion density in order to study the stability of the vacuum state against fluctuations in fermion number.

The vacuum state is defined as the ground state with lowest possible energy density amongst all other ground states which due to external influences are in states with higher energy densities.

The effective potential can be defined using $\tilde{S}_{\text{eff}}$ for spacetime independent background f fields: $V_{\text{eff}} = -\tilde{S}_{\text{eff}}/V_4$ with V_4 the volume of the spacetime box that contains the system. The stability of the vacuum state against the fluctuations in fermion number density around $\bar{\rho} = 0$ can be studied using V_{eff} , which is a canonical function of $\bar{\rho}$. Minimization of V_{eff} with respect to μ gives the vacuum value ρ_{vac} of the system since the vacuum $\bar{\rho}$ is a known function of the vacuum μ in AGCE.

The local finite density theory based upon the AGCE developed after considering the above ingredients is examined by applying it to the chiral symmetry breaking phase of the half bosonized Nambu Jona-Lasinio model for the 3+1 dimension and the chiral Gross-Neveu model for 2+1 dimensions with its effective potential given by

$$V_{\text{eff}} = -N_g \int \frac{d^D p}{(2\pi)^D} \left[\ln \left(1 + \frac{\sigma^2}{p_+^2} \right) + \ln \left(1 + \frac{\sigma^2}{p_-^2} \right) \right] + \frac{1}{4G_0} \sigma^2 + \bar{e}, \quad (7)$$

where “ D ” is the spacetime dimension, G_0 is the coupling constant and $p_\pm^2 = (p_0 \pm i\mu)^2 + \mathbf{p}^2$. It can be shown that $\partial^2 V_{\text{eff}}/\partial \mu^2|_{\mu=0} < 0$ as long as $\sigma \neq 0$ and $D \geq 2$, which implies that the $\sigma \neq 0$ state with $\bar{\rho} = 0$ is not stable against fermion number fluctuations. Such a conclusion is both theoretically and physically unacceptable.

It appears that the AGCE is not sufficient for the local finite density theory, a more general ensemble, called the asymptotic non-thermo ensemble (ANTE) is introduced to cure this pathology. The ANTE in the remote past of the system is not necessarily a strict thermal one. This is because the $\sigma \neq 0$ phase, which is called the α -phase of the massless quark system, is known to be condensed with macroscopic number of bare fermion-antifermion pairs. These pairs occupy the low energy states of the system; they block other fermions from further filling of these states. This kind of statistical blocking effect is not encoded into Eq. 7, which starts the time evolution of the system from a set of initial states having zero number of fermion-antifermion pairs that do not overlap with the states having a finite *density* of such pairs in the thermodynamic limit. Its effects can however be included in the boundary condition for the system like what has been done for the finite density cases. Therefore, it is proposed that the initial ensemble of quantum states in the ANTE for the system at zero temperature and density are those ones with negative energy states filled up to $E = -\epsilon$ rather than $E = 0$ and with positive energy states filled up to $E = \epsilon$ rather than empty. It is

expected that such a state can has sufficient overlap with the true vacuum state of the system for properly determined ϵ . In the ANTE [2], the effective potential corresponding to Eq. 7 is expressed as

$$V_{\text{eff}} = -N_g \int_{\mathcal{C}} \frac{d^D p}{(2\pi)^D} \left[\ln \left(1 + \frac{\sigma^2}{p_+^2} \right) + \ln \left(1 + \frac{\sigma^2}{p_-^2} \right) \right] + \frac{1}{4G_0} \sigma^2 + \bar{e}_{(+)} + \bar{e}_{(-)} - \bar{e}, \quad (8)$$

where $\bar{e}_{(\pm)} = 2N_g \int^{\mu_{\pm}} \frac{d^{D-1} p}{(2\pi)^{D-1}} p$. Here the upper boundary of the radial p integration in the D-1 dimensional momentum space is denoted as $\mu_{\pm} \equiv \mu \pm \epsilon$. The contour $\mathcal{C}$ for the (complex) p_0 integration is shown in Fig. 1 in which both the original Minkowski contour and the Euclidean contour are drawn. The effective potential for the $\sigma \neq 0$ case in ANTE given by the above equation has two sets of minima, the first set contains $\mu = \pm \mu_{vac}$ and $\epsilon = 0$ points, the second one includes $\mu = 0$ and $\epsilon = \pm \epsilon_{vac}$ ones with finite μ_{vac} and ϵ_{vac} . The second solutions correspond to the absolute minima, namely, the vacuum state. Although it is found that $\epsilon_{vac} < \sigma_{vac}$ in all cases that were studied, the effects of a finite vacuum ϵ can still have dynamical consequences, which is discussed in the following and elsewhere [2,1,10] due to the existence of the dark component for local observables.

The vacuum and ground states of the strong interaction could have different phases from the α -phase [11–15]. They are characterized by a condensation of diquarks. Such a possibility is interesting because it may be realized in the early universe, in astronomical objects and events, in heavy ion collisions, inside nucleons [14,16] and nuclei, etc.. For the possible scalar diquark condensation in the vacuum, a half bosonized model Lagrangian is introduced [17,2], which reads

$$\mathcal{L}_I = \frac{1}{2} \bar{\Psi} [i \not{\partial} - \sigma - i \vec{\pi} \cdot \vec{\tau} \gamma^5 O_3 - \gamma^5 \mathcal{A}_c \chi^c O_{(+)} - \gamma^5 \mathcal{A}^c \bar{\chi}_c O_{(-)}] \Psi - \frac{1}{4G_0} (\sigma^2 + \vec{\pi}^2) + \frac{1}{2G_{3'}} \bar{\chi}_c \chi^c, \quad (9)$$

where σ , $\vec{\pi}$, $\bar{\chi}_c$ and χ^c are auxiliary fields with $(\chi^c)^\dagger = -\bar{\chi}_c$ and G_0 , $G_{3'}$ are coupling constants of the model. $\mathcal{A}_c$ and $\mathcal{A}^c$ ($c = 1, 2, 3$) act on the color space of the quark; they are $\mathcal{A}_{c_1 c_2}^c = -\epsilon^{cc_1 c_2} \mathcal{A}_{c, c_1 c_2} = \epsilon^{cc_1 c_2}$ with ϵ^{abc} ($a, b = 1, 2, 3$) the total antisymmetric Levi-Civita tensor. Here $O_{(\pm)}$ are raising and lowering operators respectively in the upper and lowering 4 components of Ψ .

This model has two non-trivial phases. The vacuum expectation of σ is non-vanishing with vanishing $\chi^2 \equiv -\bar{\chi}_c \chi^c$ in the α -phase. The vacuum state in the α -phase is condensed with quark-antiquark pairs. The vacuum expectation of σ is zero with finite χ^2 that spontaneously breaks the $U(1)$ statistical gauge symmetry in the second phase, which is called the ω -phase. There is a condensation of correlated scalar diquarks and antidiquarks in the color $\bar{3}$ and 3 states in the ω -phase of the vacuum state.

Since diquarks and antidiquarks are condensed in the ω -phase, it is expected that an exchange of the role of ϵ and μ occurs. It is found to be indeed true: there are also two sets of minima for the effective potential, the first set is the one with finite $\mu = \pm \mu_{vac}$ and $\epsilon = 0$ and the second set corresponds to $\mu = 0$ and finite $\epsilon = \pm \epsilon_{vac}$ in the ω -phase of the model. But here the absolute minima of the system in the ω -phase correspond to the first set of solutions, in which the CP/T invariance and baryon number conservation are spontaneously violated due to the presence of a finite vacuum μ_{vac} . This conclusion is also applicable to the β -phase of models [11,12] with vector fermion pair and antifermion pair condensation, in which the chiral symmetry $SU(2)_L \times SU(2)_R$ is also spontaneously broken down.

The dynamics of the primary statistical gauge field are generated by $S_{\text{eff}}^{(\mu)}[\mu] = \ln Z$. Due to the statistical gauge invariance, the leading order in the derivative expansion of $S_{\text{eff}}^{(\mu)}[\mu]$ in terms of μ^α can be expressed as the following: in the normal phase, the effective action for slow varying $\mu'_\alpha = \mu_\alpha - \bar{\mu}_\alpha$ is

$$S_{\text{eff}}^{(\mu)} = \int d^4 x \left[-\frac{Z^{(\mu)}}{4} f_{\mu\nu} f^{\mu\nu} + \frac{N_g}{\pi^2} \bar{\epsilon}^2 \mu' \cdot \mu' \right] + \dots \quad (10)$$

with $f^{\alpha\beta} = \partial^\alpha \mu'^\beta - \partial^\beta \mu'^\alpha$ and $\bar{\epsilon}$ the ground state value of ϵ ; in the phase where local statistical $U(1)$ gauge symmetry is spontaneously broken down and before considering electromagnetic interaction,

$$S_{\text{eff}}^{(\mu)} = \frac{1}{2} \int d^4 x \left[i \bar{j}^\alpha \vec{j}^\beta + g^{\alpha\beta} \Pi^{(\mu)} \right] \mu'_\alpha \mu'_\beta + \dots \quad (11)$$

In both of the situations with slow varying μ'_α , $Z^{(\mu)}$ and $\Pi^{(\mu)}$, which can be extracted from the time-ordered current-current correlator $\langle T j^\alpha(x) j^\beta(x') \rangle$, are approximately constants.

The excitation corresponds to μ^α is massive for the vacuum state in the α -phase due to the statistical blocking effects. There is no long range force associated with μ^α in the α -phase. It can be shown [2] that after including the electromagnetic interaction, the spatial component of μ^α excitation is massless for the vacuum state in the β - and ω -phases in the rest frame of matter.

In summary, it is found that general local relativistic quantum field theory (in the sense of including both the zero and the finite density situations) contains dark components for local observables, statistical gauge invariance and fermionic blocking effects. The proper boundary condition for a consistent theory is that of an optimal ANTE. This view seems to be receiving support from lattice gauge theory studies [18,19]. The rich implications of the finding presented here remain to be explored.

-
- [1] S. Ying, Report No. hep-ph/9704216.
 - [2] S. Ying, Ann. Phys. (NY) **266** (1998) 295.
 - [3] See, e.g., K. Huang, “Statistical Mechanics,” Wiley, New York, 1963.
 - [4] I. M. Barbour, S. E. Morrison, E. G. Klepfish, J. B. Kogut and M. P. Lombardo, Nucl. Phys. Proc. Suppl. **A60** (1998) 220; Report No. hep-lat/9705042.
 - [5] J. Stern, Report No. hep-ph/9801282.
 - [6] A. Aspect, J. Dalibard, and G. Roger, Phys. Rev. Lett. **49** (1982) 1804.
 - [7] P. R. Tapster, J. G. Rarity and P. C. M. Owens, Phys. Rev. Lett. **73** (1994) 1923.
 - [8] W. Tittel, J. Brendel, B. Gisin, T. Herzog, H. Zbinden and N. Gisin, Phys. Rev. Lett. **81** (1998) 3563.
 - [9] G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. **23** (1998) 5039.
 - [10] S. Ying, work in progress.
 - [11] S. Ying, Phys. Lett. **B283** (1992) 341.
 - [12] S. Ying, Ann. Phys. (NY) **250** (1996) 69.
 - [13] T. Schäfer, Phys. Rev. **D57** (1998) 950.
 - [14] M. Alford, K. Rajagopal, and F. Wilczek, Phys. Lett. **B422** (1998) 247.
 - [15] R. Rapp, T. Schäfer, E. V. Shuryak, and M. Velkovsky, Phys. Rev. Lett. **81** (1998) 53.
 - [16] S. Ying, J. Phys. **G22** (1996) 293; S. Ying, Commun. Theor. Phys. **28** (1997) 301.
 - [17] S. Ying, Report No. hep-ph/9604255.
 - [18] R. Aloisio, V. Azcoiti, G. Di Carlo, A. Galante, and A. F. Grillo, Phys. Lett. **B428** (1998) 166.
 - [19] R. Aloisio, V. Azcoiti, G. Di Carlo, A. Galante, and A. F. Grillo, Report No. hep-lat/9811033.




FIG. 1. The p^0 integration contour $\mathcal{C}$ for the effective potential. The thick lines extending to positive and negative infinity represent the branch cuts of the logarithmic function. Curve “I” is the original p_0 integration contour for the theory in the Minkowski spacetime. Curve “II” corresponds to the p_0 integration contour for the theory in the Euclidean spacetime.